arXiv:1407.2983v3 [math.NT] 31 Dec 2015

A Strange Integral Equation and Some Integrals Associated with Number Theory

Alexander E Patkowski

Abstract

We utilize a combination of integral transforms, including the Laplace transform, with some classical results in analytic number theory concerning the Riemann ξ-function, to obtain a strange integral equation. We also make some other observations on a related general integral studied by N. S. Koshlyakov. We conclude with some further observations on an integral formula related to the Riemann ξ-function.

1 Introduction

In Titchmarsh [14, pg.35], we find a famous relation connecting Fourier cosine transforms with Mellin transforms

$$\int_0^\infty f(t)\lambda(t)\cos(xt)dt = \frac{1}{2i\sqrt{y}}\int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty}\phi(s-\frac{1}{2})\phi(\frac{1}{2}-s)(s-1)\Gamma(1+\frac{s}{2})\pi^{-\frac{s}{2}}\zeta(s)y^s ds, \quad (1.1)$$

where $f(t)=|\phi(it)|^2$, $\zeta(s)$ is the Riemann zeta function [6, 14], $\Gamma(s)$ is the gamma function, $\xi(s):=\frac{1}{2}s(s-1)\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})\zeta(s)$, $\lambda(t):=\xi(\frac{1}{2}+it)$ (Riemann's Ξ function), and $y=e^x$. Recall [6, 14] the famous Riemann ξ function satisfies the functional equation

$$\xi(s)=\frac{1}{2}s(s-1)\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})\zeta(s)=\frac{1}{2}s(s-1)\pi^{-\frac{1-s}{2}}\Gamma(\frac{1-s}{2})\zeta(1-s). \quad (1.2)$$

As usual, we put $\Re(s)=\sigma$, $\Im(s)=t$, for $s\in\mathbb{C}$. Many authors [4, 5, 8, 9, 12, 13, 14] have utilized equation (1.1) and its variations to obtain many interesting relations among other special functions, as well as results on $\lambda(t)$. See also [10, 11] for some interesting ideas on integrals related to Riemann's ξ function. The most widely cited example in the literature appears to be [14, eq.(2.16.1)] (for example, see [2, 7]), and its variant [14, eq.(2.16.2)]:

$$\Lambda(x):=\int_0^\infty \frac{\lambda(t)}{t^2+\frac{1}{4}}\cos(xt)dt=\frac{\pi}{2}\left(e^{x/2}-2e^{-x/2}\psi(e^{-2x})\right). \quad (1.3)$$

Here we have a slightly modified Jacobi theta function $\psi(x) = \sum_{n\geq 1} e^{-\pi n^2 x}$. As usual, we define the Laplace transform to be

$$\mathcal{L}(f)(s) := \int_0^\infty f(t)e^{-st}dt. \tag{1.4}$$

If we assume $\Re(s) > 1$, we may write

$$\begin{aligned}
&\mathcal{L}(\Lambda(x))(s-\frac{1}{2}) && (1.5)\\
&\quad = \int_0^\infty e^{-(s-\frac{1}{2})x} \int_0^\infty \frac{\lambda(t)}{t^2+\frac{1}{4}} \cos(xt)dtdx && (1.6)\\
&\quad = (s-\frac{1}{2}) \int_0^\infty \frac{\lambda(t)}{(t^2+\frac{1}{4})(t^2+(s-\frac{1}{2})^2)}dt && (1.7)\\
&\quad = \frac{\pi}{2} \int_0^\infty e^{-sx} \left(e^x - 2\psi(e^{-2x})\right) dx && (1.8)\\
&\quad = \frac{\pi}{2} \left(\frac{1}{s-1} - 2\int_0^\infty e^{-sx}\psi(e^{-2x})\right) dx && (1.9)\\
&\quad = \frac{\pi}{2} \left(\frac{1}{s-1} + 2\lim_{r\to 0^+} \int_1^r t^{s-1}\psi(t^2)dt\right) && (1.10)\\
&\quad = \frac{\pi}{2} \left(\frac{1}{s-1} - x^{-\frac{s}{2}} \sum_{n\geq 1} \lim_{r\to 0^+} \int_r^x t^{\frac{s}{2}-1} e^{-\pi n^2 t/x} dt\right) && (1.11)\\
&\quad = \frac{\pi}{2} \left(\frac{1}{s-1} - \pi^{-\frac{s}{2}} \sum_{n\geq 1} n^{-s}\gamma(\frac{s}{2}, \pi n^2)\right) && (1.12)\\
&\quad = \frac{\pi}{2} \left(\frac{1}{s-1} - \pi^{-\frac{s}{2}} \sum_{n\geq 1} n^{-s}\left(\Gamma(\frac{s}{2}) - \Gamma(\frac{s}{2}, \pi n^2)\right)\right) && (1.13)\\
&\quad = \frac{\pi}{2} \left(\frac{1}{s-1} - \frac{2\xi(s)}{s(s-1)} + \pi^{-s/2} \sum_{n\geq 1} n^{-s}\Gamma(\frac{s}{2}, \pi n^2)\right) && (1.14)
\end{aligned}$$

In the line (1.10) we made a change of variables $x = -\log t$, and in subsequent lines employed the definition of the incomplete gamma functions

$$\gamma(s,x) = \int_0^x t^{s-1}e^{-t}dt, \qquad \text{and} \qquad \Gamma(s,x) = \int_x^\infty t^{s-1}e^{-t}dt,$$

where $\Gamma(s) = \gamma(s,x) + \Gamma(s,x)$.

Note that by an instance $F(y) = e^{-ay^2}$ of the Müntz formua [14, eq.(2.11.1)], for $0 < \sigma < 1$,

$$\frac{1}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \Gamma(\frac{s}{2})\zeta(s)(\sqrt{a}x)^{-s}ds = \sum_{n\geq 1} e^{-an^2x^2} - \frac{1}{2x}\sqrt{\frac{\pi}{a}}. \tag{1.15}$$

Multiplying through by x^{v-1}, with $\Re(v) > 1$, and integrating over the interval $[0, z]$, $z > 0$, we obtain

$$\frac{z^v}{2\pi i} \int_{\sigma-i\infty}^{\sigma+i\infty} \Gamma(\frac{s}{2})\zeta(s)\frac{(\sqrt{a}z)^{-s}}{(v-s)}ds = \frac{a^{-v/2}}{2} \sum_{n\geq 1} n^{-v}\gamma(\frac{v}{2}, azn^2) - \frac{z^{v-1}}{2(v-1)}\sqrt{\frac{\pi}{a}}, \tag{1.16}$$

again for $0 < \sigma < 1$. (Note this integration is justified on the left side, since if $\Re(v-s) > 0$, then $0^{v-s} = 0$.) On the other hand the integral on the left-hand side of (1.16) is precisely $\Upsilon(v)$ defined in Theorem 1, when $a = \pi$, $z = 1$, $\sigma = \frac{1}{2}$. Hence we have arrived twice at the following result.

Theorem 1. *When $\Re(s) > 1$, we have*

$$\Upsilon(s) := (s-\frac{1}{2})\int_0^\infty \frac{\lambda(t)}{(t^2+\frac{1}{4})(t^2+(s-\frac{1}{2})^2)}dt = \frac{\pi}{2}\left(\frac{1}{s-1} - \frac{2\xi(s)}{s(s-1)} + \pi^{-s/2}\sum_{n\geq 1} n^{-s}\Gamma(\frac{s}{2},\pi n^2)\right). \tag{1.17}$$

Or equivalently, the Riemann ξ-function satisfies the integral equation

$$\int_0^\infty f(\frac{1}{2}+it)K(s,t)dt = \frac{\pi}{2}\left(\frac{-2f(s)}{s(s-1)} + g(s)\right), \tag{1.18}$$

with kernal $K(s,t) = (t^2+\frac{1}{4})^{-1}(t^2+(s-\frac{1}{2})^2)^{-1}(s-\frac{1}{2})$, where $g(s)$ may be explicitly computed for any solution f of (1.18).

The assertion that $g(s)$ may be explicitly computed follows directly from Mellin transforms in the proof of Theorem 1. From Titchmarsh [14, pg. 257] we have $\lambda(t) \ll t^A e^{-\frac{\pi}{4}t}$, which gives us

$$\Upsilon(s) \ll \int_0^\infty \frac{t^A e^{-\frac{\pi}{4}t}}{(t^2+\frac{1}{4})(t^2+(s-\frac{1}{2})^2)}dt < +\infty.$$

Clearly we have that $\Upsilon(s) = -\Upsilon(1-s)$. Upon noticing this fact, we may use (1.16) (or equivalently (1.17)) to obtain a well-known result concerning $\zeta(s)$. Computing the residue $R_{s=0} = \zeta(0)/v = -1/(2v)$ out from the integral, we may then extract the series $\frac{1}{2}\pi^{-(1-v)/2}\sum_{n\geq 1} n^{-(1-v)}\Gamma(\frac{1-v}{2},\pi n^2)$ to obtain the following expansion [1, pg.256, eq.(30)] as a direct corollary to Theorem 1:

$$\pi^{-s/2}\zeta(s)\Gamma(\frac{s}{2}) = \frac{1}{s(s-1)} + \pi^{-s/2}\sum_{n\geq 1} n^{-s}\Gamma(\frac{s}{2},\pi n^2) + \pi^{-(1-s)/2}\sum_{n\geq 1} n^{-(1-s)}\Gamma(\frac{1-s}{2},\pi n^2). \tag{1.19}$$

2 Imaginary quadratic forms and the associated integral equation

Following [6, pg.511] put

$$L_K(s,\chi) = \sum_{\mathfrak{a}} \chi(\mathfrak{a})(N\mathfrak{a})^{-s}, \tag{2.1}$$

where $\sigma > 1$, and χ maps the class group $\mathfrak{H}$ to the complex plane $\mathbb{C}$. In [8, 9], we find N. S. Koshlyakov investigating integrals related to L-functions associated with number fields as well as [12, 14]. We consider his work coupled with that of the ideas in the introduction. Let D denote a discriminant with respect to a primitive ideal $\mathfrak{a}$. Then we have the functional equation [6, eq.(22.51)]

$$\Omega_K(s,\chi) = \Omega_K(1-s,\chi), \tag{2.2}$$

where $\Omega_K(s,\chi) = (2\pi)^{-s}\Gamma(s)|D|^{\frac{s}{2}}L_K(s,\chi)$.

An important integral representation relevant to our study, which continues $L_K(s,\chi)$ to the entire complex plane, is given by Hecke [6, eq.(22.52)]:

$$\Omega_K(s,\chi) = \frac{|\mathfrak{H}|\delta(\chi)}{\bar{D}s(s-1)} + \int_1^\infty (t^{s-1}+t^{-s})\sum_{\mathfrak{a}}\chi(\mathfrak{a})e^{-2\pi t N\mathfrak{a}/\sqrt{|D|}}dt, \tag{2.3}$$

where $|\mathfrak{H}|$ is the class number, and $\bar{D}$ is 6 if $D=-3$, -4 if $D=4$, and 2 if $D<-4$. We shall prove an equivalent form of this integral representation toward the end using Theorem 2. Put $\Omega_K(\frac{1}{2}+it,\chi) = \mathfrak{O}_K(t)$, and note that

$$\int_0^\infty f(t)\mathfrak{O}_K(t)\cos(xt)dt = \frac{1}{2i\sqrt{y}}\int_{\frac{1}{2}-i\infty}^{\frac{1}{2}+i\infty}\phi(s-\frac{1}{2})\phi(\frac{1}{2}-s)(2\pi)^{-s}|D|^{s/2}\Gamma(s)L_K(s,\chi)y^s ds. \tag{2.4}$$

N.S. Koshlyakov appears to be the first to consider instances of this type of general integral with L-functions associated with number fields (see, especially, [9, pg.217–220]). We may easily evaluate the case $\phi(s)=1$, and so $f(t)=1$, for all t. Consider

$$\frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty}(2\pi)^{-s}|D|^{s/2}\Gamma(s)L_K(s,\chi)y^s ds, \tag{2.5}$$

for $c>1$, and move the line of integration to $c=\frac{1}{2}$. Since $L_K(s,\chi)$ has a simple pole at $s=1$ when χ is trivial, we compute [6, pg.512, eq.(22.50)]

$$R_{s=1} = \frac{|\mathfrak{H}|\delta(\chi)}{\bar{D}},$$

(where $\delta(\chi)=0$ when χ is non-trivial, 1 otherwise) and obtain

$$\int_0^\infty \mathfrak{O}_K(t)\cos(xt)dt = \frac{\pi}{2}\left(\frac{|\mathfrak{H}|\delta(\chi)e^{x/2}}{\bar{D}} - e^{-x/2}\Psi(e^{-x})\right), \tag{2.6}$$

where $\Psi(x) = \sum_{\mathfrak{a}}\chi(\mathfrak{a})e^{-2\pi(N\mathfrak{a})x/\sqrt{|D|}}$. (Note that (2.5) is precisely $\Psi(y^{-1})$.) Applying the same concepts as in our introduction, we may obtain the following result from (2.6).

Theorem 2. *For $\Re(s) > 1$, we have that*

$$\mathfrak{u}(s) = \int_0^\infty \mathfrak{O}_K(t)\bar{K}(s,t)dt = \frac{\pi}{2}\left(\frac{|\mathfrak{H}|\delta(\chi)}{\bar{D}(s-1)} - \Omega_K(s,\chi) + \sum_{\mathfrak{a}} \chi(\mathfrak{a})\left(\frac{\sqrt{|D|}}{2\pi N\mathfrak{a}}\right)^s \Gamma(s, \frac{2\pi N\mathfrak{a}}{\sqrt{|D|}})\right), \tag{2.7}$$

with kernel $\bar{K}(s,t) = (s-\frac{1}{2})(t^2 + (s-\frac{1}{2})^2)^{-1}$.

Due to the kernel, we again have $\mathfrak{u}(s) = -\mathfrak{u}(1-s)$, and as a direct corollary we have [6, pg. 512, eq.(22.54)]

$$\Omega_K(s,\chi) = \frac{|\mathfrak{H}|\delta(\chi)}{\bar{D}s(s-1)} + \sum_{\mathfrak{a}} \chi(\mathfrak{a})\left(\frac{\sqrt{|D|}}{2\pi N\mathfrak{a}}\right)^s \Gamma(s, \frac{2\pi N\mathfrak{a}}{\sqrt{|D|}}) + \sum_{\mathfrak{a}} \chi(\mathfrak{a})\left(\frac{\sqrt{|D|}}{2\pi N\mathfrak{a}}\right)^{1-s} \Gamma(1-s, \frac{2\pi N\mathfrak{a}}{\sqrt{|D|}}). \tag{2.8}$$

3 Other Applications and Comments for Further Research

Here we discuss some results related to (1.3) that we hope will encourage further interest. First we recall for $0 < \Re(s) < 1$, the well-known formula

$$\int_0^\infty t^{s-1}\cos(xt)dt = \frac{\Gamma(s)\cos(\frac{\pi}{2}s)}{x^s}. \tag{3.1}$$

Hence for $0 < \Re(s) < 1$,

$$\int_0^\infty t^{s-1}\left(\int_0^\infty \cos(xt)(2\psi(x^2) - \frac{1}{x})dx\right)dt = \frac{2\xi(s)\Gamma(s)\cos(\frac{\pi}{2}s)}{s(s-1)}. \tag{3.2}$$

This gives us $(0 < c < 1)$

$$2\psi(x^2) - \frac{1}{x} = \int_0^\infty \cos(xt)\left(\frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty} \frac{2\xi(s)\Gamma(s)\cos(\frac{\pi}{2}s)t^{-s}}{s(s-1)}ds\right)dt = \int_0^\infty \cos(xt)\dot{I}(t)dt, \tag{3.3}$$

say. So we may now compute (using (1.3))

$$\frac{\lambda(t)}{t^2+\frac{1}{4}} = \int_0^\infty \cos(xt)e^{-x/2}\left(\int_0^\infty \cos(e^{-x}u)\dot{I}(u)du\right)dx. \tag{3.4}$$

It is also possible to prove (3.2) using Parseval's theorem for Mellin transforms and the functional equation (1.2).

Using the work from [3] with the above observations, we consider the following result. See [3, Definition 1.1] for a criterion for an entire function to belong to the Laguerre-Pólya class.

Theorem 3. *The Riemann Hypothesis is equivalent to the statement that the integral*

$$\ddot{I}(t) := \int_0^\infty \cos(xt)e^{-x/2}\left(\int_0^\infty \cos(e^{-x}u)\dot{I}(u)du\right)dx,$$

has only real zeros. Consequently, the Riemann Hypothesis is true if and only if $\ddot{I}(t)$ *is in the Laguerre-Pólya class.*

Proof. The proof uses the integral (3.4) coupled with comparing the kernel of $\ddot{I}(t)$, given by

$$\ddot{k}(x) := e^{-x/2}\left(\int_0^\infty \cos(e^{-x}u)\dot{I}(u)du\right)dx,$$

to what is considered an 'admissible' kernel according to the definition given in [3, Definition 1.2]. For example, it is easily verified that $\ddot{k}(x)$ is even by the functional equation for the Jacobi theta function (and (3.3)), $\ddot{k}(x) > 0$ for all $x \in \mathbb{R}$, but $\frac{d}{dx}\ddot{k}(x) > 0$ for all $x > 0$. Hence the kernel is monotone increasing for $x > 0$. The fact that the kernel satisfies these properties tells us that $\ddot{I}(t)$ is a real entire function, and it is possible to have only real zeros. The remainder of the proof utilizes the well-known criterion that the Riemann hypothesis is equivalent to the statement that all the zeros of $\lambda(t)$ are real, together with [3, Definition 1.1]. □

At this point we are left with some questions regarding the solutions of Theorem 1. First, in light of equation (1.18) resembling a "modified" Fredholm integral of the second kind, does the integral equation in Theorem 1 admit application of a kind of Fredholm theory? Can anything new be said by attempting a Neumann series solution?

More interesting relations may be obtained by expanding $L_K(s,\chi)$ in (2.4) into a sum of Epstein zeta functions, with formulas related to the work in [8, 9, 15]. This would allow us to obtain integrals related to a theta functions of the form $\sum_{n,m\in\mathbb{Z}} \chi_{n,m} q^{an^2+bnm+cm^2}$, $4ac - b^2 > 0$,where $q = e^{-\pi x}$, $x > 0$. It may also be of interest to look at the integral $\ddot{I}(t)$ by explicitly computing $\dot{I}(u)$ using the Residue theorem.

References

[1] J.M. Borwein D. M. Bradley, R. E. Crandall, *Computational strategies for the Riemann zeta function,* Journal of Computational and Applied Mathematics 121 (2000) 247–296.

[2] J. B. Conrey and A. Ghosh, *Turn inequalities and zeros of Dirichlet series associated with certain cusp forms,* Trans. Amer. Math. Soc. 342 (1994), 407-419

[3] G. Csordas, *Fourier Transforms of Positive Definite Kernels and the Riemann ξ-function,* Computational Methods and Function Theory, Volume 15, Issue 3, pp 373–391 (2015).

[4] A. Dixit, *Series transformations and integrals involving the Riemann Ξ-function,* J. Math. Anal. Appl., 368, (2010), 358–373.

[5] A. Dixit *Character analogues of Ramanujan type integrals involving the Riemann Ξ-function,* Pacific J. Math., 255, No. 2 (2012), 317–348

[6] H. Iwaniec and E. Kowalski, *Analytic number theory,* American Mathematical Society Colloquium Publications, vol. 53, American Mathematical Society, Providence, RI, 2004.

[7] H. Ki, *The Riemann Ξ-function under repeated differentiation,* J. of Number Theory, Volume 120, Issue 1, September 2006, Pages 120–131.

[8] N. S. Koshlyakov, *Investigation of some questions of the analytic theory of a rational and quadratic field. I* Izv. Akad. Nauk SSSR Ser. Mat., 18:2 (1954), 113–144.

[9] N. S. Koshlyakov, *Investigation of some questions of the analytic theory of a rational and quadratic field. II* (in Russian), Izv. Akad. Nauk SSSR Ser. Mat. 18 No. 3, 213–260 (1954).

[10] A. Kuznetsov, *Integral representations for the Dirichlet L-functions and their expansions in Meixner–Pollaczek polynomials and rising factorials,* Integral Transforms and Special Functions 18 (2007), No. 11-12, 809–817.

[11] A. Kuznetsov, *Expansion of the Riemann Ξ function in Meixner-Pollaczek polynomials,* Canadian Math. Bulletin 51 (2008), No. 4, 561–569.

[12] S. Ramanujan, *New expressions for Riemann's functions $\xi(s)$ and $\Xi(t)$,* Quart. J. Math., 46: 253–260, 1915.

[13] R. Spira, *The Integral Representation for the the Riemann Ξ function,* J. of Number Theory, Volume 3, Issue 4, November 1971, Pages 498–501

[14] E. C. Titchmarsh, *The theory of the Riemann zeta function,* Oxford University Press, 2nd edition, 1986.

[15] K. S. Williams and Zhang Nan-Yue, *On the Epstein zeta function,* Tamkang J. Math. 26 (1995), 165-176.

1390 Bumps River Rd.

Centerville, MA 02632

USA

E-mail: alexpatk@hotmail.com